

MODULE 11: INTIMACY POST-STROKE

Name: _____ Date: _____

1. What 2 non-physical post-stroke effects also affect intimacy post-stroke? (2 points)

2. Can resumption of sexual intimacy lead to another stroke? Yes or No (1 point)

3. The PLISSIT model provides a base for addressing intimacy post-stroke with stroke survivors. Describe the PLISSIT model. (4 points)

Quiz

- You may use your binder as a reference to answer these questions
- Submit your completed quiz to the Nurse Clinician or designate for marking
- Your test will be returned to you to keep in your binder

TRUE/FALSE Questions

(CIRCLE the correct letter)

- T** **F** 1. A stroke survivor's most common fear in relation to resumption of sexual activity is that it can cause another stroke.
- T** **F** 2. Less than 50% of stroke survivors experience a decrease in sexual activity.
- T** **F** 3. Medications that are prescribed to treat stroke risk factors can affect a stroke survivor's sexual desire and performance.
- T** **F** 4. Stroke survivors are not concerned about intimacy post-stroke, as their focus is solely on their physical recovery.
- T** **F** 5. Healthcare providers do not routinely provide stroke survivors with education and support regarding intimacy post-stroke because they are uncomfortable and feel they lack the knowledge to do so.
- T** **F** 6. Psychological rather than physical factors account for most sexual disruption post-stroke.
- T** **F** 7. Engaging in intimacy post-stroke is normal and healthy for stroke survivors.

SCORE: _____ / **14**

Stroke Unit/Medical Unit Nurse Clinician/Designate

SIGNATURE