

MODULE 4: DIAGNOSTICS AND ASSESSMENTS



Learning Objectives

Upon completion of this module, nurses will be able to:

- Describe the acute stroke nursing assessment
- Understand why neurological assessments and functional assessments are important
- State the standardized neurological assessments
- Explain the relevant stroke investigations and why they are being done



4.1 Acute Care Nursing Assessment

Acute stroke patients require skilled and knowledgeable professionals attending to their care. Organized stroke units create the opportunity to develop a critical mass that facilitates the development of expertise of clinicians. Expertise in stroke assessment by clinicians ensures that stroke patients are offered appropriate investigations and receive appropriate and timely care.

“The nurse is often the first to see changes and early warning signs that may predict a neurological crisis. Familiarity with a stroke nurse assessment promotes the nurse’s ability to gather accurate patient information, identify stroke emergencies and promote timely referrals to appropriate specialists and time-dependent investigations” (HSFO, Faaast FAQs, 2007, page 9).



A stroke assessment includes:

(HSFO, Faaast FAQs, 2007, p. 10)

1. ABC

Begin a general assessment of the moment you first encounter the person. Take note any signs of acute distress that are present and assess ABC (airway-breathing-circulation).

2. Health History

The purpose of the health history is to collect subjective data and combine it with objective data from the physical examination. The combined database is then used to make a judgment or diagnosis about the health status of the individual (Jarvis, 2000).

3. Vital Signs

Temperature, heart rate, blood pressure, respiration rate, pain assessment and oxygen saturation are all measured and monitored according to Standards of Care for the unit. The frequency of monitoring vital signs is outlined in Module 3: Pre-hospital and Emergency Management.

4. Pupils

Record the size of the pupils in mm using the pupil scale prior to the application of the light stimulus. Indicate the reaction of pupils as either:

+ = Brisk Reaction **S** = Sluggish **-** = No Reaction

If the eyes are closed due to swelling, record “**C**”

***It is critical to report a change in either pupil.**

5. Standardized Neurological Assessment

A neurological (neuro) assessment provides a standardized method to rapidly identify emerging stroke complications, and will provide a better patient prognosis.

Early identification of emerging stroke complications may:

- Lead to early intervention
- Limit the extension of neurological damage
- Impact patient outcomes

4.2 Standardized Neurological Assessments (CNS, NIHSS, GCS)



The use of standardized and validated stroke assessment tools in both the acute and rehabilitation settings enables **sound decision making** and **care planning**.

What are you looking for?

Symptoms of change in neurological status

- Restlessness
- Combativeness
- Confusion
- Severe headache
- Lethargy
- Decline in motor strength
- Decrease in coordination
- Change in balance
- Change in speech/language

(HSFO, Faaast FAQs, 2007)

There are three neurological assessment tools available for nurses: Canadian Neurological Scale (CNS), the National Institutes of Health Stroke Scale (NIHSS) and the Glasgow Coma Scale (GCS) (HSFO, 2003; RNAO, 2005). The CNS and the NIHSS are assessments specific to stroke. You can find the training for these two in *Module 5: Canadian Neurological Scale (CNS) Training*, or *Module 5: National Institute for Health Stroke Scale (NIHSS) Training*, depending on which tool your organization uses.

Canadian Neurological Scale* (CNS):

The Canadian Neurological Scale is an assessment tool for evaluating and monitoring the neurological status of acute stroke patients. It has been found to be brief, valid and reliable, and can be administered in approximately 5 minutes.

The CNS assesses:

- level of consciousness (LOC)
- Orientation
- Speech
- Motor function of the face, arms and legs

A limitation of the scale may be that some useful measures are omitted. For example, it does not include assessment of cerebellar or brainstem function (HSFO, 2003; RNAO, 2005).

*For detailed instructions on how to perform these assessments refer to Module 5 in this ASU Resource

National Institute of Health Stroke Scale* (NIHSS):

The National Institute of Health Stroke Scale was designed for use in both clinical trials and everyday practice to assess stroke severity and degree of recovery. The NIHSS has demonstrated reliability and validity for patients with stroke. It can take up to 20 minutes to administer until you are quite comfortable with each item. Then, it may be completed in 5-10 minutes.

The NIH Stroke Scale (NIHSS) assesses:

- LOC
- Visual fields
- Motor response
- Sensation
- Language
- Neglect

It is an **11 item scale** that measures various physiological deficits associated with stroke:

- LOC
- Best gaze
- Visual field testing
- Facial paresis
- Arm motor function
- Leg motor function
- Limb ataxia
- Sensory
- Best language
- Dysarthria
- Extinction and inattention

*For detailed instructions on how to perform these assessments refer to Module 5 in this ASU Resource

The NIHSS allows us to:

- Objectively quantify our clinical exam (give it a score)
- Determine if the patient's neurological status is improving or deteriorating
- Provide standardization in assessment (from one nurse to the next – the same exam is performed)
- Communicate patient status to staff (nurses and *Physicians* talking the same language)
- Aid in decision making re: tPA treatment, acute care needs, needs at discharge

A limitation of the scale is that it has a low sensitivity to small changes.

(HSFO, 2003; RNAO, 2005)

Glasgow Coma Scale (GCS):

The Glasgow Coma Scale is a standardized and valid neurological assessment tool for assessing level of consciousness or coma.

It is a neurological assessment that is widely used by the neurological and neurosurgery community and is found in the curriculum of most undergraduate nursing programs.

It lacks specificity and applicability when applied to stroke patients, as most do not have impaired LOC.

***Where do I document these neurological assessments in my organization?
*Follow your organization's charting guidelines.***

4.3 Other Assessments



There are many more assessments relevant to stroke inpatients. Some of these assessments are administered by other members of the team. Consult with your Nurse Clinician or Stroke Nurse to learn more about any one of the assessments listed below.

- **Function:** AlphaFIM® (an abbreviated version of the FIM® Instrument to assess function and disability in the acute care setting)
- **Pain:** Visual Analog Scale, Wong-Baker Faces Rating Scale, Numeric Rating Scale
- **Skin Breakdown:** Braden Risk Assessment
- **Balance:** Berg Balance Scale
- **Cognition:** Montreal Cognitive Assessment (MoCA), Mini Mental
- **Depression:** Geriatric Depression Scale, Patient Health Questionnaire -9 (PHQ9), Hospital and Anxiety Depression Scale (HADS)
- **Deep Vein Thrombosis:** Follow organization's DVT prophylaxis guidelines
- **Swallowing:** Toronto Bedside Swallowing Screening Test (TOR BSST), etc.
- **Continence**
- **Nutrition/Hydration**
- **Oral Care:** Oral Health Assessment Tool (OHAT)
- **Sleep Apnea:** Refer to 2012 Canadian Best Practice Recommendations for Stroke Care (2.8.1): Screening and Assessment for Sleep Apnea
- **Driving:** Refer to 2013 Canadian Best Practice Recommendations (5.11): Life Roles and Activities – Return to Driving

4.4 Investigations

Refer to your organizations pre-printed orders for specific details regarding investigations.

CT – HEAD is ordered STAT in emergency department to rule out brain hemorrhage or identify other etiologies. The CT scan may appear normal for several hours after the onset of stroke. In ischemic stroke, a larger infarct may appear as a hypodense area (core) surrounded by area of intermediate density (penumbra). Hypodense area is most commonly seen in 24-48 hrs. It is necessary to repeat in 2 days. CT does not show posterior circulation strokes well.

CT – Angiogram (CTA) (Head and Neck) examines blood flow through the arteries in the neck and brain – dye is injected into a vein and a series of rapid-image x-rays are taken as the dye travels through.

*Metformin is usually held 48hrs post CT with contrast.

All patients with suspected acute stroke, presenting within acute stroke treatment time windows, must undergo immediate noncontrast brain CT imaging, and vascular imaging with CTA including extracranial and intracranial arteries to guide hyperacute care (2015 Canadian Best Practice Recommendations for Stroke Care: Hyperacute). CT and CTA should be done back to back.

Multiphase CTA in Acute Stroke

Multi-phase CTA provides cerebral angiograms in three phases (peak arterial, peak venous and late venous) after the injection of contrast.

Good collateral flow is associated with:

- Better clinical outcomes
- Less chance of haemorrhagic transformation
- A greater possibility of successful recanalization

Delaney, F., et al. (n.d.)

Multiphase CTA in Acute Stroke provides a better indication of collateral flow and is essential to help guide patient selection for EVT.

Imaging criteria influencing EVT selection:

- Presence of a proximal occluded artery, which is a target lesion amenable to EVT
- Presence of good collaterals on multiphase CTA
- Small to moderate ischemic core as defined by an ASPECTS score of >6 on noncontrast CT.

Stotts, G. & Krings, T. (2016).

Magnetic Resonance Imaging (MRI)/Magnetic Resonance Angiography (MRA) may be ordered if CT is normal, may detect ischemic changes, smaller infarcts and TIAs presenting acutely.

Carotid Doppler is an ultrasound of carotid arteries to detect blockages.

Blood tests confirm (or refute) suspicions of specific clinical conditions:

- **Complete Blood Count (CBC)** is used to evaluate overall health and detect a wide range of disorders including anemia, infection, etc.
- **Levels of cardiac enzymes** (including **troponin** and **creatinine kinase**) can confirm heart muscle damage and extent of damage.
- **Creatinine** findings can show reduced renal blood flow, atherosclerosis.
- **Glucose Random Testing** indicates blood glucose levels in the blood.
- **PTT/INR** Partial Thromboplastin Time indicates the function of all coagulation factors. International Normalized Ratio (INR) is a ratio used to monitor the effectiveness of blood thinning drugs such as warfarin (Coumadin).
- **Lipid Profile** is a complete fasting lipoprotein profile that will show total blood cholesterol level, HDL, LDL and triglyceride levels.
- **HbA1c** is a test that measures the amount of glycated hemoglobin in your blood. Glycated hemoglobin is a substance in red blood cells that is formed when glucose attaches to hemoglobin. It indicates how well blood sugar has been controlled over the previous 2-3 months. HbA1c results can contribute to a diagnosis of diabetes.

Chest X-RAY will determine whether the heart is enlarged or if fluid is accumulating in the lungs.

Echocardiogram (ECHO) uses ultrasound to show the shape, texture and movement of heart valves, as well as the size of heart chambers and how well they are working. An ECHO may be done to diagnose heart valve problems or identify a cardiac cause of stroke.

Electrocardiogram (ECG) is a test to examine a graphic record of the heart's electrical impulses, determining timing and duration of each electrical phase in the heart beat. The result will determine if a heart attack has occurred and monitors for changes in heart rhythm.

Holter Hook up/Telemetry Monitoring (24 and 48 hr) is a portable and remote, respectively, monitor of the patient's cardiac rhythm. They may be used to detect heart rhythm abnormalities during daily activities, ruling out atrial fibrillation as a cause of stroke.

O₂ saturation monitoring picks up presence of hypoxia which may exacerbate and worsen ischemic damage. If hypoxic, possible causes of respiratory compromise (pneumonia, partial airway obstruction, hypoventilation, and atelectasis) must be considered and managed.

Optional:

Transesophageal echocardiogram (TEE) is an invasive procedure to obtain a more detailed view of the heart structures from the esophagus.

Electroencephalogram (EEG) uses electrodes on the patient's scalp to investigate electrical activity in the brain to rule out seizure.

References

Casaubon LK, Boulanger J-M, Blacquiére D, Boucher S, Brown K, Goddard T, Gordon J, Horton M, Lalonde J, LaRivière C, Lavoie P, Leslie P, McNeill J, Menon B K, Moses B, Penn M, Perry J, Sneider E, Tymianski D, Foley N, Smith E, Gubitz G, Hill M. D, Glasser E, and Lindsay P on behalf of the Heart and Stroke Foundation of Canada, Canadian Stroke Best Practices Advisory Committee, (2015). Canadian Stroke Best Practice Recommendations: Hyperacute Stroke Care Guidelines, Update 2015. *International Journal of Stroke* Volume 10, Issue 6, p. Pages 924–940

Coutts S, Kelloway L, on behalf of the Prevention of Stroke Writing Group. *Chapter 2: Prevention of Stroke*. In Lindsay MP, Gubitz G, Bayley M, and Phillips S (Editors) on behalf of the Canadian Stroke Best Practices and Standards Working Group. *Canadian Best Practice Recommendations for Stroke Care: 2012*; Ottawa, Ontario Canada: Canadian Stroke Network.

Dawson D, Knox J, McClure A, Foley N, and Teasell R, on behalf of the Stroke Rehabilitation Writing Group. *Chapter 5: Stroke Rehabilitation*. In Lindsay MP, Gubitz G, Bayley M, and Phillips S (Editors) on behalf of the Canadian Stroke Best Practices and Standards Advisory Committee. *Canadian Best Practice Recommendations for Stroke Care: 2013*; Ottawa, Ontario Canada: Heart and Stroke Foundation and the Canadian Stroke Network.

Delaney, F., et al. (n.d.). Multiphase CT Angiography in Acute Ischaemic Stroke. Retrieved from <https://radiopaedia.org/articles/multiphase-ct-angiography-in-acute-ischaemic-stroke>

Eskes G, Salter K, on behalf of the Mood and Cognition in Stroke Writing Group and the Evidence-Based Review in Stroke Rehabilitation Team. *Chapter 7: Mood and Cognition in Stroke*. In Lindsay MP, Gubitz G, Bayley M, and Phillips S (Editors) on behalf of the Canadian Stroke Best Practices and Standards Advisory Committee. *Canadian Best Practice Recommendations for Stroke Care: 2013*; Ottawa, Ontario Canada: Canadian Stroke Network.

Heart and Stroke Foundation of Ontario, (2003). *Heart and Stroke Best Practice Guidelines for Stroke Care*. A resource for implementing optimal stroke care. Toronto: Heart and Stroke Foundation of Ontario.

Heart and Stroke Foundation of Ontario and Canadian Stroke Network. (2007). *Faaast FAQs for Stroke Nurses*. Ottawa: Heart and Stroke Foundation of Canada.

Heart and Stroke Foundation of Canada and Canadian Stroke Network, (2007/2008). *Acute Stroke Management Resource*. (Power point presentations). Retrieved from http://www.heartandstroke.on.ca/site/c.pvI3leNWJwE/b.5346923/k.8D25/HCP__English.htm.

Registered Nurses' Association of Ontario, Nursing Best Practice Guidelines. (2005). *Stroke Assessment Across the Continuum of Care*. Toronto: Registered Nurses' Association of Ontario.

Schlegel, D., Kolb,SJ., and Kasner, SE. (2003). *Utility of the NIH Stroke Scale as a Predictor of Hospital Disposition*. *Stroke*. 34, 134-137.

Stotts, G. & Krings, T. (2016). *Endovascular Treatment for Acute Ischemic Stroke* [PowerPoint Presentation] Feb 3rd and 5th.

Web Sources

- www.labtestsonline.org
- <http://www.strokingengine.ca/>
- www.strokebestpractices.ca