

ACUTE STROKE UNIT ORIENTATION

2023

MODULE 5: SWALLOWING, NUTRITION AND ORAL CARE

Name: ______ Date: _____

- 1. What estimates range of acute stroke patients have some degree of dysphagia?
 - a. Approximately 5-25%
 - b. Approximately 25-52%
 - c. Approximately 29-67%
 - d. Approximately 80-90%
- 2. Acute stroke patients are to remain NPO until swallowing ability has been determined. These patients:
 - 1. May have ice chips
 - 2. May have po meds with sips of water
 - 3. May have nothing by mouth
 - 4. May have thickened fluids until further assessed by the Speech-Language Pathologist.

Choose Answer:

- a. 1 and 2
- b. 1, 2 and 3
- c. 3
- d. 4

Quiz

- You may use the resource as a reference to answer these questions.
- Submit your completed quiz to the Nurse Clinician or designate for marking.

3.	 Dysphagia is the a. Loss or impaired ability to chew and/or swallow b. Inability to comprehend spoken language c. Loss of expressive oral ability d. Motor planning impairment about the mouth
4.	Complications associated with dysphagia may include: a. Malnutrition and dehydration b. Reduced quality of life c. Aspiration pneumonia and airway obstruction d. All of the above
5.	During a normal swallow, the food is chewed and mixed to form a bolus during the: a. Pharyngeal phase b. Esophageal phase c. Oral phase d. Peristalsis phase
6.	During a <i>normal swallow,</i> the pharyngeal phase is a. Voluntary b. Involuntary/reflexive c. A physiological response to aspiration d. Often accompanied by coughing
7. 1.	List 5 common signs that your stroke patient is having difficulty swallowing. (5 points)

What <i>may</i> (1 point)	not occur whe	n a patient as _i	pirates silently?	

- 10. Proper oral care MUST INCLUDE:
 - a. Entire mucosa
 - b. Sulci (spaces between the cheeks and gums)
 - c. Upper and lower teeth
 - d. All of the above

TRUE/FALSE Questions

(CIRCLE the correct letter)

Τ	F	1.	Patients on thickened fluids are at increased risk the risk for dehydration.				
Т	F	2.	Feeding a patient while they are lying down is okay as long as they are comfortable.				
Т	F	3.	All stroke survivors should be screened for dysphagia as soon as they are awake and alert and before ANY oral intake is allowed.				
Т	F	4.	Presence of a gag reflex does not protect against aspiration.				
SCORE: / 18							
Stroke Unit/Medical Unit Nurse Clinician/Designate							

SIGNATURE